

# **Solar and Wind Energy Resource Assessment (SWERA)**

## **High Resolution Solar Radiation Assessment for China**

**Final Report for a Country Assessment prepared by** 

**Christoph Schillings<sup>1</sup>**

**Richard Meyer<sup>2</sup>**

**Franz Trieb<sup>1</sup>**

<sup>1</sup> Deutsches Zentrum für Luft- und Raumfahrt, DLR-Stuttgart  
Institut für Technische Thermodynamik, Pfaffenwaldring 38-40, D-70569 Stuttgart, Germany

<sup>2</sup> Deutsches Zentrum für Luft- und Raumfahrt, DLR-Oberpfaffenhofen  
Institut für Physik der Atmosphäre, D-82234 Weßling, Germany

submitted to UNEP / GEF

October 2004

## Content

1. Method description (satellite data, *GHI*-method, *DNI*-method)
2. Model output (*GHI*, *DNI*)
3. Comparison with ground measurements (if available)
4. References

## Notice

This report was prepared for the SWERA project funded by GEF / UNEP. The results presented here were produced by state-of-the-art methods with best-known data. The DLR (Deutsches Zentrum für Luft- und Raumfahrt e.V.) does not make any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the DLR.

This report is electronically available at <http://swera.unep.net>

Contact the responsible author:


Christoph Schillings  
DLR - Deutsches Zentrum für Luft- und Raumfahrt, e.V.  
Pfaffenwaldring 38-40  
D - 70569 Stuttgart

tel     ++49(0)711/6862-784  
fax     ++49(0)711/6862-783  
mail    christoph.schillings@dlr.de  
www    <http://www.dlr.de/system>

## **1 Method description**

### **Satellite Data**

The high resolution solar radiation assessment is based on data of the geostationary satellite Meteosat. Due to the location of the participating SWERA countries, data of Meteosat 7 (M-7) for the years 2000, 2001 and 2002 (for Ghana, Kenya and Ethiopia) and data of Meteosat 5 (M-5) for the years 2000, 2002 and 2003 (for Bangladesh, West-China, Nepal and Sri Lanka) are used. M-5 has its position at 0° latitude and 63° East longitude, M-7 is located at an orbit at 0° latitude and 0° longitude. Figure 1 gives the field of view of both satellites which scans the specific area every 30 minutes with a spatial resolution of 5x5 km<sup>2</sup>.



**Figure 1:** The solar irradiance data is derived from Meteosat at 0° (red circle) and at 63° East (orange circle). The brightened area marks the quantitatively analyzable region. (Meyer et al., 2004).

Data of the visible (VIS) channel, which gives the reflection of the system earth/atmosphere (including clouds) and data of the infrared (IR) channel, which represents the temperature of the surface and atmosphere, are used for gathering information about the clouds. Both are used in a different way to assess the global horizontal (*GHI*) and the direct normal radiation (*DNI*) at ground. Additionally, data of the most important atmospheric components that attenuate the radiation, namely ozone, water vapor and aerosols, are used to take into account the clear-sky conditions of the atmosphere. In the following, the method for deriving *DNI* based on the DLR method and the method for deriving *GHI*, based on a combined method of DLR and SUNY, is described.

### **Method for Direct Normal Radiation (*DNI*)**

The calculation of *DNI* is based on the clear-sky model of Bird and Hulstrom (1981) as described in Iqbal (1983) which was modified by Schillings et al. (2004) for taking into account cloudy conditions with

$$DNI = 0.9751 \cdot I_0 \cdot \tau_R \cdot \tau_{Gas} \cdot \tau_{Ozon} \cdot \tau_{WV} \cdot \tau_{Ae} \cdot \tau_{vis} \cdot \tau_{ir} \quad (1)$$

Each atmospheric transmittance coefficient  $\tau_{cl}$  is calculated separately using atmospheric input data. All equations for calculating the clear-sky transmittances are described in Iqbal (1983).

Transmittance for Rayleigh scattering

$$\tau_R = \exp\left[-0.0903m_a^{0.84}\left(1.0 + am_p - am_p^{1.01}\right)\right] \quad (2)$$

Transmittance for equally distributed gas (mainly O<sub>2</sub> and CO<sub>2</sub>)

$$\tau_{Gas} = \exp\left(-0.0127am_p^{0.26}\right) \quad (3)$$

Transmittance for ozone

$$\tau_{Ozon} = 1 - \alpha_{Ozon} \quad (4)$$

$$\alpha_{Ozone} = 0.1611\chi(1.0 + 139.48\chi)^{-0.3035} - 0.002715\chi(1.0 + 0.044\chi + 0.0003\chi^2)^{-1} \quad (5)$$

$\chi = u \cdot am$ , with the vertical ozone layer thickness  $u$  in cm[NTP] and the airmass  $am$ .

Transmittance for water vapor

$$\tau_{WV} = 1 - \alpha_{WV} \quad (6)$$

$$\alpha_{WV} = 2.4959\gamma\left[(1.0 + 79.034\gamma)^{0.6828} + 6.385\gamma\right]^{-1} \quad (7)$$

$\gamma = w \cdot am$ , with the pressure-corrected relative optical path length of precipitable water  $w$  in cm[NTP].

Transmittance for aerosols

$$\tau_{Ae} = \exp\left[-k_a^{0.873}\left(1.0 + k_a - k_a^{0.7088}\right)am_p^{0.9108}\right] \quad (8)$$

$$k_a = 0.2758k_{a\lambda=0.38\mu m} + 0.35k_{a\lambda=0.5\mu m} \quad (9)$$

with the aerosol optical thickness  $k_{a\lambda}$  at the wavelength 0.38  $\mu m$  und 0.5  $\mu m$ .

Transmittance for clouds

using the visible Cloud-Index  $CI_{vis}$

$$\tau_{vis} = e^{(-CI_{vis} \cdot 0.1)} \quad (10)$$

and using the infrared Cloud-Index  $CI_{ir}$

$$\tau_{ir} = e^{(-CI_{ir} \cdot 0.07)} \quad (11)$$

For the clear-sky atmospheric transmittance, the air mass is needed, which is calculated by

$$am = \frac{1}{\left[\cos\Theta_Z + 0.15(93.885 - \Theta_Z)\right]^{-1.253}} \quad (12)$$

The adaption for actual ground elevation is made by

$$am_p = am \cdot \frac{p}{1013.25} \quad (13)$$

with

$$\frac{p}{p_0} = \exp(-0.0001184z) \quad (14)$$

The clear-sky radiation is calculated each 20 minutes (10, 30, 50 minutes of each hour) for the maps and each 5 minutes (5,10,15,...,55,60 minutes each hour) for the time series. The influence of the clouds is taken into account based on hourly averages calculated from half hourly Meteosat data. Therefore all maps (monthly and annual average daily sums) and time series are based on an hourly calculation of the radiation. The DLR -model output for *DNI* is sampled at a 10km spatial resolution.

### Method for Global Horizontal Radiation (*GHI*)

The calculation of *GHI* is based on the method of Perez et al (2002):

$$GHI = ktm \cdot G_{hc} \cdot (0.0001 \cdot ktm \cdot G_{hc} + 0.9) \quad (15)$$

with *ktm*

$$ktm = 2.36 \cdot CI^5 - 6.2 \cdot CI^4 + 6.22 \cdot CI^3 - 2.63 \cdot CI^2 - 0.58 \cdot CI + 1 \quad (16)$$

*GHI* is calculated using the cloud information based on infrared (IR) and visible (VIS) Meteosat data following the procedure of Schillings et al. (2004), which lead to a single Cloud-Index *CI* with

$$CI = \max(CI_{vis}, CI_{ir}) \quad (17)$$

For the determination of the clear-sky global irradiance  $G_{hc}$  the new formulation as described in Perez et al (2002) is used with

$$G_{hc} = cg1 \cdot I_0 \cdot \cos \Theta_z \cdot \exp(-cg2 \cdot am \cdot (fh1 + fh2 \cdot (TL - 1))) \exp(0.01 \cdot am^{1.8}) \quad (18)$$

with

$$\begin{aligned} cg1 &= (0.0000509 \cdot alt + 0.868) \\ cg2 &= (0.0000392 \cdot alt + 0.0387) \\ I_0 &= \text{Solar constant (eccentricity corrected)} \\ \Theta_z &= \text{solar zenith angle} \\ fh1 &= \exp(-alt / 8000) \\ fh2 &= \exp(-alt / 1250) \\ am &= \text{elevation corrected air mass} \\ alt &= \text{altitude in meters} \\ T_L &= \text{Linke turbidity} \end{aligned}$$

Instead of using the Linke turbidity  $T_L$  for the parameterization of the clear-sky atmosphere, data of the atmospheric components ozone, water vapor and aerosols are used to calculate it. As these atmospheric data are also used for the *DNI* results are more consistent. To derive  $T_L$  from atmospheric data we use the following formula as described by Ineichen and Perez (2002) with

$$TL = ((11.1 \cdot \ln(b \cdot \frac{I_0}{B_{ncl}})) / am) + 1 \quad (19)$$

$$\text{with } b = 0.664 + (0.163 / fh1) \quad (20)$$

and the clear-sky direct normal irradiance  $B_{ncl}$

$$B_{ncl} = I_0 \cdot \tau_{ra} \cdot \tau_{ae} \cdot \tau_{o3} \cdot \tau_{ga} \cdot \tau_{wv} \quad (21)$$


The calculation of transmittance coefficients  $\tau_i$  and the used atmospheric input data are described above in the section describing the method to retrieve *DNI*.

The clear-sky radiation is calculated each 20 minutes (10, 30, 50 minutes of each hour) for the maps and each 5 minutes (5,10,15,...,55,60 minutes each hour) for the time series. The influence of clouds is taken into account hourly, therefore all maps (monthly and annual average daily sums) and time series are based on an hourly calculation of the radiation. The DLR/SUNY-model output for *GHI* is sampled at a 10km spatial resolution.

## Input Data

### Elevation


For the air mass elevation adaption, the elevation from the GLOBE database from the USGS U.S. Geological Survey [[http://rockyweb.cr.usgs.gov/elevation/dpi\\_dem.html](http://rockyweb.cr.usgs.gov/elevation/dpi_dem.html)] is used, (Hastings and Dunbar, 1998).



**Figure 2:** Elevation from GLOBE.

### Ozone


The monthly ozone data are taken from TOMS as published by the NASA/GSFC TOMS Ozone Processing Team [<http://toms.gsfc.nasa.gov/>], (McPeters et al., 1998).



**Figure 3:** Ozone monthly average for February 2003 in [DU] from TOMS

### Water vapor


The daily water vapor data are taken from the NOAA-CIRES Climate Diagnostics Center in Boulder Colorado, USA (NCEP/NCAR) [<http://www.cdc.noaa.gov/>] (Kalnay et al., 1996).



**Figure 4:** Water vapor daily mean for 7. February 2003 in cm[NTP] from NCEP/ NCAR-Reanalysis

### Aerosol

The monthly climatological aerosol optical thickness data are taken from NASA-GACP, [<http://gacp.giss.nasa.gov/index.html>], (Mishchenko et al, 2002).



**Figure 5:** Aerosol optical thickness for February from NASA-GACP.

### Clouds

The hourly cloud information is based on half-hourly Meteosat-5 IR and VIS data (EUMETSAT, © 2004). The determination of the cloud indices is described in detail in Mannstein et al. (1999) and Schillings et al. (2004). The basic approach for deriving VIS cloud information is described with

$$CI_{vis} = \frac{\rho - \rho_{min}}{\rho_{max} - \rho_{min}} \quad (22)$$

where  $\rho$  is the actual reflectivity measured by the satellite,  $\rho_{min}$  corresponds to the surface albedo and  $\rho_{max}$  is the maximum reflectivity measured for overcast cloudy conditions. The similar approach is used for IR-data, with the actual, minimum and maximum brightness temperatures  $T$  measured by the satellite:

$$CI_{-ir} = \frac{T_{\min} - T}{T_{\min} - T_{\max}} \quad (23)$$




**Figure 6:** Field of view of Meteosat (M-7, IR-channel) © 2004, EUMETSAT.

## **2 Model output**


The solar radiation is calculated for the complete country for the years 2000, 2001 and 2002. The data are made available in a digital GIS-format (ESRI Vector-Shapefile). Within this report, maps of the annual average daily total sum of *GHI* and *DNI* are presented. The complete database (ESRI-Shapefile and MS-Access database) can be downloaded from the SWERA-homepage (<http://swera.unep.net>). Within the ESRI Vector-Shapefile, 3 annual and 36 monthly average daily total sums of *GHI* and *DNI* are given for each 10km x 10km geo-referenced pixel as shown in the following figures. Additional, hourly time series for the same time period for several interesting sites are delivered in a separate ASCII-File.

**Important notice:** The following maps show classified values of *kWh/m<sup>2</sup>/day* with a common color ramp for all SWERA countries. This is done to give a first impression of the solar regime for each country and for easier comparison with other countries. The provided digital GIS data (available at <http://swera.unep.net>) give the real (and not classified!) values in *Wh/m<sup>2</sup>/day* for each geo-referenced pixel with a signal resolution of 1 *Wh/m<sup>2</sup>/day*.


## Global Horizontal Radiation



**Figure 7:** Annual average daily total sum of *GHI* kWh/m<sup>2</sup>/day for West-China 2000




**Figure 8:** Annual average daily total sum of *GHI* in kWh/m<sup>2</sup>/day for West-China 2002




**Figure 9:** Annual average daily total sum of GHI in kWh/m<sup>2</sup>/day for China 2003


### Direct Normal Radiation



**Figure 10:** Annual average daily total sum of *DNI* in kWh/m<sup>2</sup>/day for West-China 2000



**Figure 11:** Annual average daily total sum of  $DNI$  in  $\text{kWh/m}^2/\text{day}$  for West-China 2002



**Figure 12:** Annual average daily total sum of *DNI* in kWh/m<sup>2</sup>/day for West-China 2003

### **3 Comparison with ground measurement in China**

No comparison was performed due to missing ground measurement data.

#### **Acknowledgement:**

All Meteosat data is under copyright of EUMETSAT (© 2004), Darmstadt, Germany. Many thanks for access to the data set go to the crew of the MARF (Meteosat Archive and Retrieval Facility, Darmstadt) and to our colleagues from DLR-DFD (Deutsches Fernerkundungs-Datenzentrum). Data on aerosol is provided by NASA-GACP Global Aerosol Climatology Project. We also acknowledge the use of the water vapor data from the NCEP Reanalysis by NOAA-CIRES Climate Diagnostics Center, Boulder, Colorado, US and the use of the TOMS ozone data provided by the NASA-Goddard Space Flight Center (GSFC), Washington, DC, USA. Special thanks to Rüdiger Buell and Hermann Mannstein at the Institut für Physik der Atmosphäre for doing the archive logistics and cloud data processing of the Meteosat 5 and Meteosat 7 data.

### **4 References**

- Bird R.E. and R.L. Hulstrom (1981): Review, evaluation and improvement of direct irradiance models. *J. of Sol. Energy Engineering* **103**, 182-192.
- Hastings D.A. and P.K. Dunbar (1998): Development & Assessment of the Global Land One-km Base Elevation Digital Elevation Model (GLOBE). *International Society of Photogrammetry and Remote Sensing (ISPRS) Archives* **32** (4), 218-221. (<http://www.ngdc.noaa.gov/seg/topo/globe.shtml>)
- Ineichen P. and R. Perez (2002): A new airmass independent formulation for the Linke turbidity coefficient. *Solar Energy*, **73** (3), 151-157.
- Iqbal M. (1983): An introduction to solar radiation. Academic Press, Toronto.
- Kalnay E., M. Kanamitsu, R. Kistler, W. Collins, D. Deaven, L. Gandin, M. Iredell, S. Saha, G. White, J. Woollen, Y. Zhu, A. Leetmaa, R. Reynolds, M. Chelliah, W. Ebisuzaki, W. Higgins, J. Janowiak, K.C. Mo, C. Ropelewski, J. Wang, R. Jenne and D. Joseph (1996): The NMC/NCAR 40-Year Reanalysis Project. *Bulletin of the American Meteorological Society* **77** (3), 437-472.
- Mannstein H., H. Broesamle, C. Schillings and F. Trieb (1999): Using a METEOSAT cloud index to model the performance of solar-thermal power stations. *Procs. of Eumetsat Conference Copenhagen*, 239-246.
- McPeters R.D., P. K. Bhartia, A. J. Krueger, J. R. Herman, C. G. Wellemeyer, C. J. Seftor, G. Jaross, O. Torres, L. Moy, G. Labow, W. Byerly, S. L. Taylor, T. Swissler and R. P. Cebula (1998): Earth Probe Total Ozone Mapping Spectrometer (TOMS) Data Products User's Guide. *NASA Technical Publication* 1998-206895.
- Meyer R., C. Hoyer, E. Diedrich, C. Schillings, S. Lohmann, M. Schroedter-Homscheidt, R. Buell and F. Trieb (2004): Solar Energy Mining: High-Resolution Meteosat-based Service for Solar Radiation Products for Europe, Africa and Asia, Proc. of the EuroSun 2004 Conference, Freiburg im Breisgau, Germany, June 20-23 2004, vol. 3, 857-863.

Mishchenko M., J.E. Penner and D. Anderson (Ed.) (2002): Global Aerosol Climatology Project. Special Issue of Global Aerosol Climate, *J. of Atmospheric Science* **59** (3).

Perez R., P. Ineichen, K. Moore, M. Kmiecik, C. Chain, R. George and F. Vignola (2002): A new operational satellite-to-irradiance model - description and validation. *Solar Energy*, **73** (5), 307-317.

Schillings C., H. Mannstein and R. Meyer (2004): Operational method for deriving high resolution direct normal irradiance from satellite data. *Solar Energy* **76**, 475-484.